

Science[Prev](#) | [Table of Contents](#) | [Next](#)

www.sciencemag.org

Science 25 July 1997:

Vol. 277 no. 5325 pp. 500–504

DOI: 10.1126/science.277.5325.500

• ARTICLES

Biotic Control over the Functioning of EcosystemsF. Stuart Chapin III^{*}, Brian H. Walker, Richard J. Hobbs, David U. Hooper, John H. Lawton, Osvaldo E. Sala, David Tilman

☐ Author Affiliations

F. S. Chapin III and D. U. Hooper are in the Department of Integrative Biology, University of California, Berkeley, CA 94720–3140, USA. B. H. Walker is in the Division of Wildlife and Ecology, Commonwealth Scientific and Industrial Research Organisation (CSIRO), Post Office Box 84, Lyneham, ACT 2602, Canberra, Australia. R. J. Hobbs is in the Division of Wildlife and Ecology, CSIRO, LMB 4 Post Office Midland, Western Australia 6056, Australia. J. H. Lawton is at the National Environmental Research Council Centre for Population Biology, Imperial College, Silwood Park, Ascot SL5 7PY, UK. O. E. Sala is on the Faculty of Agronomy, University of Buenos Aires, Avenida San Martin 4453, Buenos Aires 1417, Argentina. D. Tilman is in the Department of Ecology, Evolution, and Behavior, University of Minnesota, St. Paul, MN 55108, USA.


ABSTRACT

Changes in the abundance of species—especially those that influence water and nutrient dynamics, trophic interactions, or disturbance regime—affect the structure and functioning of ecosystems. Diversity is also functionally important, both because it increases the probability of including species that have strong ecosystem effects and because it can increase the efficiency of resource use. Differences in environmental sensitivity among functionally similar species give stability to ecosystem processes, whereas differences in sensitivity among functionally different species make ecosystems more vulnerable to change. Current global environmental changes that affect species composition and diversity are therefore profoundly altering the functioning of the biosphere.

The two most dramatic ecological trends of the past century are human-induced changes in biotic diversity and alterations to the structure and functioning of ecosystems (1). Ecosystem processes, such as productivity, nitrogen mineralization rate, and nitrate leaching, respond directly to human modification of ecosystems and to changes in atmospheric composition and climate (2). Changes in biota result from habitat conversion and land use change, reducing genetic and species diversity; and from the introduction of exotic species, leading to a homogenization of the global biota (1). In addition to the ethical, aesthetic, and economic concerns raised by this situation, we contend that these biotic changes will influence ecosystem processes sufficiently to alter the future state of the world's ecosystems and the services they provide to humanity. The current global extinction rate, which is 100 to 1000 times greater than prehuman levels (3, 4), and the loss of local diversity due to management practices have the potential to affect ecosystem processes strongly on both local and global scales. This article focuses on species, but there are also functionally important differences at other scales (5). For example, genetic diversity in crops improves resistance to disease and reduces the risk of large yield losses caused by epidemics (6). At large scales, the diversity and patterning of ecosystems in a landscape affect regional processes, such as nutrient transport from terrestrial to aquatic ecosystems across riparian zones (7).

Species differ in the rates and pathways by which they process resources, in their effects on the physical environment, and in their interactions with other species. Thus, changes in species composition are likely to alter ecosystem processes through changes in the functional traits of biota (Fig. 1). This can alter ecosystem processes, such as nitrogen uptake by vegetation, which in turn

modifies community processes such as competition and herbivory, feeding back to further changes in community composition. Species-induced changes in ecosystem processes can also alter regional processes (Fig. 1) such as methane emissions from beaver ponds (8) or nutrient transfers to aquatic ecosystems (7), extending the impacts beyond the original zone of species change. Some species-induced changes in ecosystem and regional processes alter ecosystem services to humans, such as purity of water supplies (9) and forest productivity (10). Species diversity can influence these same processes (11), for two reasons. First, the number of species in a community is a measure of the probability of the presence of species with particularly important traits; second, greater diversity allows a greater range of traits to be represented in the ecosystem, providing opportunities for more efficient resource use in a variable environment.



Changes in species composition and diversity will affect the functioning of ecosystems most strongly when species differ in their effects on ecosystem processes or in their response to environmental change. In the first case, by definition a change in species composition or abundance must affect ecosystem functioning. In the second case, differential environmental sensitivity among functionally similar species gives stability (resistance and resilience) to ecosystem processes, whereas differences in sensitivity among functionally different species make ecosystems more vulnerable to change.

Species Traits

What species traits are most likely to affect ecosystem processes? Traits with profound effects are those that (i) modify the availability, capture, and use of soil resources such as water and nutrients, (ii) affect the feeding relationships (trophic structure) within a community, and (iii) influence the frequency, severity, and extent of disturbances such as fire (12).

Resource dynamics. The supplies of water, nutrients, and space are important “bottom-up” controls of ecosystem structure and dynamics (13), so the introduction or loss of species that alter resource availability strongly affects ecosystem processes. A species could alter resource supply by tapping an otherwise unused source; examples are the introduction of mycorrhizal fungi into mine tailings, where they enhance phosphorus availability to their plant hosts (14); the introduction of nitrogen fixers in systems where they did not previously occur (15); or the introduction of deep-rooted species, such as *Tamarix* (salt cedar) into deserts (16) or *Eucalyptus* into mediterranean ecosystems (17), that tap previously inaccessible water and nutrient sources. These species shifts can have major implications for runoff to lakes and streams, salinization of reservoirs, or recovery of degraded ecosystems (18).

Organisms modify rates of element and energy transfer within ecosystems in species-specific ways (19). For example, there are substantial differences among plant species in litter quality (20), effects on soil temperature and moisture (21), and exudation of organic compounds from roots (22), all of which affect nutrient mineralization. Animals influence the resource base of the ecosystem by changing the distribution (23) or importation of nutrients; an example is nutrient import from oceans to streams by migratory salmon. Similarly, termites transport clay particles from subsoils to near the surface, increasing water and nutrient retention (24). Microbial functional groups that mediate nutrient cycling, including heterotrophs, nitrifiers, and denitrifiers, differ in their effects on nutrient availability and loss from ecosystems. However, less is known about how the composition of microbial communities within these broad guilds affects processes at the ecosystem level (25); there may be substantial overlap in function at the level of microbial species. Although nitrogen and phosphorus dynamics are sensitive to changes in microbial community composition and biomass, carbon cycling may be less so (26). In aquatic systems, groups of microbes of different sizes differ in their effects on nutrient and energy flow; small-bodied taxa retain nutrients and energy within the microbial community, whereas the larger plankton are consumed by grazers large enough to provide food to fish (27). Changes in microbial biota will have major ramifications for ecosystems in which only one or a few species fill a functional group, in which species-specific interactions control ecosystem dynamics (for example, mutualisms or pathogens), or in which microbes that are “strong controllers” are sensitive to disturbance (25).

Widespread changes in species composition that alter resource use can have regional-scale impacts. Simulations suggest that conversion of the Amazon basin from forest to pasture would cause a permanent warming and drying of South America, because the shallower roots of grasses access less water, leading to less evapotranspiration and greater energy dissipation as sensible heat (28).

Trophic structure. Some of the most dramatic changes in ecosystem processes have resulted from the

introduction or loss of predators or diseases that have large “keystone” effects, which are effects that are substantially greater than would be expected from the biomass of the species (9, 29). For example, the introduction of exotic fish can radically alter the abundance of other fish or insects that eat zooplankton, which in turn graze on algae, the abundance of which determines water quality and use by people (9,29). Removal by humans of elephants or other keystone mammalian herbivores leads to encroachment of woody plants into savannas (30). Such changes in the abundance of keystone herbivores may have contributed to past shifts in the distribution of biomes (31), which, in turn, influence climate (32). Microbial trophic dynamics in both soil and aquatic systems can also have large effects on the turnover and fate of nutrients. In soil, the grazing of microorganisms by protozoans often leads to higher rates of nutrient turnover and greater nitrogen and phosphorus availability to plants (33). Similarly, epidemic diseases, such as rinderpest in Africa, act as keystone species by modifying competitive interactions and community structure (34). A species that is innocuous in its home environment frequently becomes invasive in a new location, if introduced without its herbivores, diseases, and other natural control agents (35).

Disturbance regime. Animals or plants that alter the disturbance regime increase the importance of nonequilibrium processes, such as colonization, relative to equilibrium processes, such as competition (5, 19). For example, gophers and pigs disturb the soil, creating sites for seedling establishment and favoring early successional, short-lived species (36). Beavers in North America are “ecosystem engineers” (19) that alter hydrology, aeration, and carbon inputs to soil, influencing the production of greenhouse gases such as methane and CO₂(8). Plants can reduce disturbance rates by stabilizing soils and reducing wind and soil erosion. Thus, even species that are uncommon in mid- and late succession can be critical to the long-term sustainability of an ecosystem (12). On the other hand, introduction of grasses into forest or shrubland ecosystems can increase the frequency of fire and cause a replacement of forest by savanna (37). Disturbances created by overgrazing can alter the albedo of the land surface and change regional patterns of temperature and precipitation. In one case, the reduced heating and convective uplift of the overlying air mass caused less advection of moisture from the Mediterranean and reduced precipitation (38). The increased drought amplified the regional reduction in biomass and production. In another case, the reduction in transpiration resulting from overgrazing in northern Mexico increased sensible heat flux, causing regional warming (39).

Indirect species effects. Species that by themselves have small effects on ecosystem processes can have large indirect effects if they influence the abundance of species with large direct ecosystem effects. Thus, a seed disperser or pollinator that has little direct effect on ecosystem processes may be essential for the persistence of a canopy species that has greater direct ecosystem impact (40). Unknown indirect effects are often cited by ecologists as justification for the importance of species diversity. However, there is currently no theoretical framework to predict when these indirect effects are most important.

Species Differences in Environmental Response

Species differences in response to environmental change can either provide stability or trigger dramatic functional changes, depending on the traits of the species involved. Species that are similar to one another in their effects on ecosystem processes but differ in their response to the environment provide stability, because any decrease in the abundance of one species will be compensated for by increases in other functionally similar species (41, 42). For example, in response to acidification of a lake, the biomass of various groups such as cladocerans, copepods, rotifers, and total zooplankton remained high despite the loss of component species from each group, owing to compensatory increases by other taxa (43). The more functionally similar species there are in a community—that is, the greater the diversity within a functional group—the greater will be its resilience in responding to environmental change, if those species differ in environmental responses (41, 42). For example, because of the presence of drought-tolerant species, diverse grasslands maintained higher productivity in response to drought than did grasslands whose diversity had been reduced by experimental addition of nutrients (44). In Western Australia, functionally similar *Acacia* species have different temperature thresholds for germination, and therefore different species colonize after fires of different intensities (45), thus ensuring the replacement of nitrogen after fire across a broad range of burn conditions. Conversely, the fewer species there are in any assemblage, the more likely it is that extinctions will alter ecosystem processes associated with that functional group (46). For example, overhunting of sea otters, the only major predator of sea urchins in the western Pacific, increased the abundance of urchins, which grazed down kelp and eliminated kelp forests over extensive areas (47). As time scales increase, an ecosystem will experience a wider range of conditions, increasing the importance of diversity among functionally similar species. Thus, genetic and species diversity per se are important to long-term maintenance of community and ecosystem structure and processes. This argues that no two species are ecologically redundant, even if they appear similar in their ecosystem effects under one particular set of environmental conditions.

In contrast to the buffering provided by ecologically similar species, species that differ in their response to the environment and in their effects on ecosystem processes can make ecosystems vulnerable to change. For example, rising concentrations of atmospheric CO₂ can reduce plant transpiration, resulting in increased magnitude or duration of soil moisture (48). This, in turn, can shift the competitive balance from grasses to shrubs, promoting shrub encroachment into grasslands and savannas and causing replacement of one biome by another. If extensive, shrub encroachment

could reduce regional albedo and enhance regional warming. Similarly, model simulations suggest that forest expansion into tundra at the arctic treeline could have accounted for up to 50% of the high-latitude climate warming that occurred during the Holocene warming 6000 years before the present (49).

Impacts of Species Number

Recent experimental and observational studies reveal that biotic diversity can affect multiple ecosystem processes (5). In all cases, this dependence comes from diversity being a summary variable that measures the extent of differences in the traits represented in the ecosystem. Studies in Minnesota grasslands (50), greenhouses (51), and controlled environment chambers (52) show that either increased plant diversity, or the types of species represented at different levels of diversity, led to increased plant productivity. Other studies, including those with mixed-species agriculture, find that mixture yields are less than yields of the most productive monoculture but often exceed the average productivity of monocultures of the component species (5, 53, 54). In the Minnesota grassland study, greater plant diversity increased the uptake of limiting soil nitrogen and reduced leaching loss of nitrogen, which could, in the long term, help maintain soil fertility. This may occur because species have complementary patterns of resource acquisition or because higher diversity increases the probability of the presence of productive species (11, 54).

Higher diversity might increase the stability of ecosystem processes in at least three ways (5, 41, 44). First, a high diversity of trophic interactions in diverse ecosystems might provide alternative pathways of energy flow and therefore more stable energy flow among trophic levels (5, 55). Second, higher species diversity might reduce the susceptibility of ecosystems to invasion by species with novel ecosystem effects after disturbance (56). Third, higher diversity can reduce the spread of plant pathogens by increasing the average distance between individuals of a given species (53). For each process, there are simple, but mostly untested, mechanisms that cause the rates and intensities of various ecosystem processes to depend on species diversity.

However, in comparisons among natural ecosystems, there is often no clear relation between species diversity and function; for example, the high resistance of diverse Australian shrub communities to plant invasion (57) and high susceptibility to introduced pathogens (58) compares with the vulnerability of diverse South African shrublands to plant invasion (59). This suggests that factors other than diversity are also important and that we still have much to learn about the role of high levels of natural diversity. This highlights the need for carefully designed comparative and experimental studies of systems with high diversity (60).

Future Scenarios

Given the frequently strong effects of species composition and diversity on ecosystem processes seen in experimental studies, what can we expect in the future? Land use change currently has the largest effect on biodiversity (Fig. 2), but changes in atmospheric composition and climate will likely have increasing impacts. The current rapid rates of deforestation, urbanization, and overexploitation, whether through overgrazing or overfishing (61), strongly affect species composition and diversity. Areas in which there is substantial nitrogen deposition also exhibit reduced diversity (62). The effects of altered land use on diversity may depend on changes in atmospheric composition and climate. For example, habitat fragmentation may prevent species migration in response to climatic change (32), thus causing greater species loss than would either factor by itself.



View larger version:
[In this page](#) [In a new window](#)
[Download PowerPoint Slide for Teaching](#)

Figure 2

Many human activities drastically simplify the species richness of ecosystems. (A) shows an ancient hay meadow on the Lower Derwent Ings in North Yorkshire, England. “Ing” is a Viking word meaning flooded meadow. The Ings flood every winter, remain damp in the summer, receive no fertilizer or pesticide inputs, and are cropped for hay and grazed in traditional land use practices that have their origins in Viking Britain. The resulting seminatural ecosystems are extremely rich in plants, insects, and birds. They are also very easily destroyed and drastically simplified (B) by draining, ploughing, the elimination of winter flooding, and conversion to arable agriculture. Similar changes in land use are severely altering the biological diversity of ecosystems all over the world.

General circulation models predict geographically variable changes in climate, with temperature increases being most pronounced at high latitudes and precipitation changing with more complex patterns (63). Because high-latitude ecosystems have low diversity, climatically mediated changes in species composition will likely have large ecosystem effects in these areas. Land use change will also be heterogeneous because population growth, demand for food, and the suitability of ecosystems to produce that food vary regionally (64). The combination of rapid land use change and high diversity in the tropics have made these areas particularly vulnerable to species loss, with largely unknown functional consequences. In regions where changes in both land use and climate are strong, such as in midcontinent regions, we expect to see the greatest changes in species composition, diversity, and

ecosystem processes. Often areas that are hot spots for diversity, such as riparian corridors, are the areas that experience the greatest human impact and have the largest effects on landscape processes.

Recent research on biotic controls on ecosystem processes has only scratched the surface of the complex web of interactions that govern the functioning of the biosphere. Comparative studies, both manipulative and observational, are required to determine why some ecosystems are more resistant and resilient to interference than others.

Conclusions

Both types of species present and diversity per se have important influences on ecosystem processes and services. The apparent conflict between the perspectives that each species is important (65) and that there is ecological redundancy among species (42) is resolved when biotic composition is considered in terms of functional types of organisms and their environmental responses. Changes in the abundance of species that differ in ecosystem consequences should affect process rates or patterns, whereas the abundance of species with similar ecological effects should give stability (resistance and resilience) to ecosystems in the face of increasingly rapid human-induced environmental change. Loss of a keystone species or of all species in a major functional group will, by definition, have large ecosystem effects. Efforts to identify and protect such species and groups often yield demonstrable near-term benefits. Of increasing concern is the loss of species that have similar ecosystem effects but differ in their environmental responses. Loss of such species may reduce ecosystem resilience and the capacity to adjust to ever-increasing rates of environmental change. This latter role of diversity is not adequately represented in current international conventions, but it may be one of the most important mechanisms by which we sustain the long-term functioning of ecosystems and the services they provide to society.

Species effects on ecosystems occur at all scales, from local to global, and their effects may be intense or subtle. Ecologists are only now beginning to establish the theoretical, empirical, and experimental frameworks to understand and predict how changes in species composition affect ecosystem processes. In a world in which local and global species extinctions are accelerating and exotic species are entering communities at unprecedented rates, links between species and ecosystem processes are emerging as a problem of fundamental concern.

_* To whom correspondence should be addressed. E-mail fschapin@socrates.berkeley.edu

REFERENCES AND NOTES

1. P. M. Vitousek, H. A. Mooney, J. Lubchenco, J. M. Melillo, *Science* **277**, 494 (1997). [Abstract/FREE Full Text](#)
2. P. M. Vitousek, *Ecology* **75**, 1861 (1994). [CrossRef](#) [Web of Science](#)
3. S. I. Dimm, C. I. Russell, J. L. Gittleman, T. M. Brooks, *Science* **269**, 347 (1995). [Abstract/FREE Full Text](#)
4. J. H. Lawton and R. M. May, Eds., *Extinction Rates* (Oxford Univ. Press, Oxford, 1995).
5. K. G. Johnson, K. A. Vogt, H. J. Clark, O. J. Schmitz, D. J. Vogt, *Trends Ecol. Evol.* **11**, 372 (1996).
6. J. E. Vanderplank, *Disease Resistance in Plants* (Academic Press, Orlando, FL, 1984).
7. W. T. Peterjohn, D. L. Correll, *Ecology* **65**, 1466 (1984). [CrossRef](#) [Web of Science](#)
8. S. D. Bridham, C. A. Johnston, J. Pastor, K. Updegraff, *Bioscience* **45**, 262 (1995). [CrossRef](#) [Web of Science](#)
9. S. P. Carpenter and J. F. Kitchell, Eds., *The Trophic Cascade in Lakes* (Cambridge Univ. Press, Cambridge, 1993).
10. D. Binkley, P. Collins, R. Bell, D. Sachs, D. Myrold, *Ecology* **73**, 2022 (1992). [CrossRef](#) [Web of Science](#)
11. R. M. Baker, S. W. Pacala, E. A. Bazzaz, C. D. Canham, S. A. Levin, *Global Change Biol.* **1**, 373 (1995); D. Tilman, C. Lehman, K. Thomson, *Proc. Natl. Acad. Sci. U.S.A.* **94**, 1857 (1997). [Abstract/FREE Full Text](#)
12. P. M. Vitousek, *Oikos* **57**, 7 (1990); F. S. Chapin III, J. Lubchenco, H. I. Reynolds, in *Global Biodiversity Assessment*, V. H. Heywood, Ed. (Cambridge Univ. Press, Cambridge, 1995), pp. 23–45.
13. H. Jenny, *The Soil Resources: Origin and Behavior* (Springer-Verlag, New York, 1980); F. S. Chapin III, M. S. Torn, M. Tateno, *Am. Nat.* **148**, 1016 (1996). [CrossRef](#) [Web of Science](#)
14. M. F. Allen, *The Ecology of Mycorrhizae* (Cambridge Univ. Press, Cambridge, 1991).
15. P. M. Vitousek, I. P. Walker, I. D. Whiteaker, D. Mueller-Dombois, P. A. Matson, *Science* **238**, 802 (1987). [Abstract/FREE Full Text](#)
16. T. E. A. van Hylckama, *Geol. Surv. Prof. Pap.* 491-E (1974).
17. M. Robles and F. S. Chapin III, *Madroño* **42**, 349 (1995).
18. D. I. McFarlane, R. I. George, P. Farrington, in *Towards Sustainable Production and Nature Conservation*, P. I. Hobbs and D. A. Saunders, Eds. (Springer-Verlag, New York, 1992), pp. 146–186; D. A. Perry, M. P. Amaranthus, I. G. Borchers, S. L. Borchers, R. E. Brainerd, *Bioscience* **39**, 230 (1989). [CrossRef](#) [Web of Science](#)
19. J. H. Lawton and C. G. Jones, in *Linking Species and Ecosystems*, C. G. Jones and J. H. Lawton, Eds. (Chapman & Hall, New York, 1995), pp. 141–150.

20. R. J. Scholes and B. H. Walker, *An African Savanna* (Cambridge Univ. Press, Cambridge, 1993).
21. S. F. Hobbie, in *Arctic and Alpine Biodiversity: Patterns, Causes and Ecosystem Consequences*, F. S. Chapin III and C. Körner, Eds. (Springer-Verlag, Berlin, 1995), pp. 213–224.
22. D. R. Zak *et al.*, *Plant Soil* **151**, 105 (1993). [CrossRef](#) [Web of Science](#)
23. C. O. Rateli, P. C. White, S. E. Marlean Jr., E. A. Ditalka, R. D. Collier, in *An Arctic Ecosystem: The Coastal Tundra at Barrow, Alaska*, I. Brown, P. C. Miller, I. I. Tieszen, F. L. Bunnell, Eds. (Dowden, Hutchinson and Ross, Stroudsburg, PA, 1980), pp. 335–410.
24. M. Lepage, *J. Anim. Ecol.* **53**, 107 (1984).
25. D. U. Hooper, D. I. Hawksworth, S. S. Dhillon, in *Global Biodiversity Assessment*, V. H. Heywood, Ed. (Cambridge Univ. Press, Cambridge, 1995), pp. 433–443.
26. E. Kandeler, C. Kampichler, O. Horack, *Biol. Fertil. Soils* **23**, 299 (1996).
27. I. P. Bormann, in *Comparative Analyses of Ecosystems: Patterns, Mechanisms and Theories*, J. Cole, G. Lovett, S. Findlay, Eds. (Springer-Verlag, New York, 1991), pp. 97–119.
28. J. Shukla, C. Nobre, P. Sellers, *Science* **247**, 1322 (1990). [Abstract/FREE Full Text](#)
29. M. E. Power *et al.*, *Bioscience* **46**, 609 (1996). [CrossRef](#) [Web of Science](#)
30. P. N. Owen-Smith, *Megaherbivores: the Influence of Very Large Body Size on Ecology* (Cambridge Univ. Press, Cambridge, 1988); J. B. Wilson and D. Q. Agnew, *Adv. Ecol. Res.* **23**, 263 (1992).
31. S. A. Zimov *et al.*, *Am. Nat.* **146**, 765 (1995). [CrossRef](#) [Web of Science](#)
32. J. M. Malillo, I. C. Prentice, C. D. Farouhar, E.-D. Schulze, O. E. Sala, in *Climate Change 1995: The Science of Climate Change*, J. T. Houghton *et al.*, Eds. (Cambridge Univ. Press, Cambridge, 1996), vol. 2, pp. 445–481.
33. R. E. Ingham, J. A. Trofymow, E. R. Ingham, D. C. Coleman, *Ecol. Monogr.* **55**, 119 (1985).
34. W. I. Bond, in *Ecosystem Function and Biodiversity*, E.-D. Schulze and H. A. Mooney, Eds. (Springer-Verlag, Berlin, 1993), pp. 237–253.
35. M. J. Crawley, *Philos. Trans. R. Soc. London Ser. B* **314**, 711 (1986). [Abstract/FREE Full Text](#)
36. P. I. Hobbie, H. A. Mooney, *Ecology* **72**, 59 (1991); [CrossRef](#) [Web of Science](#); P. M. Kotanen, *Ecography* **18**, 190 (1995).
37. C. M. D'Antonio and P. M. Vitousek, *Annu. Rev. Ecol. Syst.* **23**, 63 (1992). [Web of Science](#)
38. J. G. Charney, W. I. Ojirk, S.-H. Chow, J. Kornfield, *J. Atmos. Sci.* **34**, 1366 (1977). [Search Google Scholar](#)
39. R. C. Balling, *Clim. Change* **13**, 99 (1988). [Search Google Scholar](#)
40. R. T. Paine, *J. Anim. Ecol.* **49**, 667 (1980); I. H. Cushman, in *Islands: Biological Diversity and Ecosystem Function*, P. M. Vitousek, L. L. Loope, H. Adersen, Eds. (Springer-Verlag, Berlin, 1995), pp. 135–147.
41. S. I. McNair, *Am. Nat.* **111**, 515 (1977); [CrossRef](#) [Web of Science](#); F. S. Chapin III, G. R. Shaver, *Ecology* **66**, 564 (1985). [CrossRef](#) [Web of Science](#)
42. B. H. Walker, *Concepts Biol.* **6**, 18 (1992); *ibid.* **9**, 747 (1995); I. H. Lawton and V. K. Brown, in *Biodiversity and Ecosystem Function*, E.-D. Schulze and H. A. Mooney, Eds. (Springer-Verlag, Berlin, 1993), pp. 255–270.
43. T. M. Frost, S. P. Carpenter, A. P. Ives, T. K. Kratz, in *Linking Species and Ecosystems*, C. G. Jones and J. H. Lawton, Eds. (Chapman & Hall, New York, 1995), pp. 224–239.
44. D. Tilman, J. A. Downing, *Nature* **367**, 363 (1994). [CrossRef](#) [Web of Science](#)
45. L. Atkins and R. J. Hobbs, *CALM Science(Suppl.)* **4**, 67 (1995).
46. C. S. Holling, in *Sustainable Development and the Biosphere*, W. C. Clark and R. E. Munn, Eds. (Cambridge Univ. Press, Cambridge, 1986), pp. 292–317.
47. J. A. Estes, J. F. Palmisano, *Science* **185**, 1058 (1974). [Abstract/FREE Full Text](#)
48. C. E. Owensby *et al.*, in *Carbon Dioxide and Terrestrial Ecosystems*, G. W. Koch and H. A. Mooney, Eds. (Academic Press, San Diego, CA, 1996), pp. 147–162.
49. J. A. Foley, J. E. Kutzbach, M. T. Coe, S. Levis, *Nature* **371**, 52 (1994). [CrossRef](#)
50. D. Tilman, D. Wedin, J. Knops, *ibid.* **379**, 718 (1996). [Search Google Scholar](#)
51. S. Naeem, I. E. Thompson, S. P. Lawler, I. H. Lawton, P. M. Woodfin, *Philos. Trans. R. Soc. London Ser. B* **347**, 249 (1995). [Abstract/FREE Full Text](#)
52. —, *Nature* **368**, 734 (1994). [CrossRef](#) [Web of Science](#)
53. J. H. Vandermeer, in *Agroecology*, C. P. Carroll, J. H. Vandermeer, P. M. Rosset, Eds. (McGraw-Hill, New York, 1990), pp. 481–516; B. R. Trenbath, *Adv. Agron.* **26**, 177 (1974).
54. D. U. Hooper, *Ecology*, in press; D. U. Hooper and P. M. Vitousek, *Ecol. Monogr.*, in press.
55. R. H. MacArthur, *Ecology* **36**, 533 (1955). [CrossRef](#) [Web of Science](#)
56. C. S. Elton, *The Ecology of Invasions by Animals and Plants* (Methuen, London, 1958); D. Tilman, *Ecology* **78**, 81 (1997). [CrossRef](#) [Web of Science](#)
57. R. J. Hobbs and L. Atkins, *Aust. J. Ecol.* **13**, 171 (1988).
58. R. T. Wills, *ibid.* **18**, 145 (1993).
59. D. M. Richardson and P. M. Cowling, in *Fire in South African Mountain Fynbos: Ecosystem Community and Species Response at Swarthskloof*, R. W. van Wilgen, D. M. Richardson, F. J. Kruger, H. J. van Hensbergen, Eds. (Springer-Verlag, Berlin, 1992), pp. 161–181.
60. P. I. Hobbie, D. M. Richardson, C. W. Davis, in *Mediterranean-Type Ecosystems: The Function of Biodiversity*, G. W. Davis and D. M. Richardson, Eds. (Springer-Verlag, New York, 1995), pp. 1–42.
61. J. F. Richards, in *The Earth as Transformed by Human Actions*, B. L. Turner II, Ed. (Cambridge

- Univ. Press, Cambridge, 1993), pp. 163–178.
62. F. Berendse and R. Aerts, *Funct. Ecol.* **1**, 293 (1987).
63. A. Kattenberg *et al.* in *Climate Change 1995: The Science of Climate Change*, J. T. Houghton *et al.*, Eds. (Cambridge Univ. Press, Cambridge, 1996), pp. 285–357.
64. J. Alcamo, Ed. *IMAGE 2.0: Integrated Modeling of Global Climate Change* (Kluwer Academic, Dordrecht, Netherlands, 1994).
65. P. P. Ehrlich and A. H. Ehrlich, *Extinction: The Causes and Consequences of the Disappearance of Species* (Random House, New York, 1981).

THIS ARTICLE HAS BEEN CITED BY OTHER ARTICLES:

MURICID DRILLING PREDATION AT HIGH LATITUDES: INSIGHTS FROM THE SOUTHERNMOST ATLANTIC

PALAIOS 1 January 2013: 33–41.

[Abstract](#) [Full Text](#) [Full Text \(PDF\)](#)

Foliar stoichiometry under different mycorrhizal types in relation to temperature and precipitation in grassland

J Plant Ecol 7 December 2012: rts042v1–rts042.

[Abstract](#) [Full Text](#) [Full Text \(PDF\)](#)

Biodiversity and Ecosystem Function

Science 13 January 2012: 174–175.

[Abstract](#) [Full Text](#) [Full Text \(PDF\)](#)

Diversity increases biomass production for trematode parasites in snails

Proc R Soc B 7 December 2008: 2707–2714.

[Abstract](#) [Full Text](#) [Full Text \(PDF\)](#)

From the Cover: Opposing plant community responses to warming with and without herbivores

Proc. Natl. Acad. Sci. USA 26 August 2008: 12353–12358.

[Abstract](#) [Full Text](#) [Full Text \(PDF\)](#)

Effects of Predator Hunting Mode on Grassland Ecosystem Function

Science 15 February 2008: 952–954.

[Abstract](#) [Full Text](#) [Full Text \(PDF\)](#)

Parallel Declines in Pollinators and Insect–Pollinated Plants in Britain and the Netherlands

Science 21 July 2006: 351–354.

[Abstract](#) [Full Text](#) [Full Text \(PDF\)](#)

A microtiter plate procedure for evaluating fungal functional diversity on nitrogen substrates

Mycologia 1 March 2006: 353–363.

[Abstract](#) [Full Text](#) [Full Text \(PDF\)](#)

The Compact City Fallacy

Journal of Planning Education and Research 1 September 2005: 11–26.

[Abstract](#) [Full Text \(PDF\)](#)

Polar Bears in a Warming Climate

Integr. Comp. Biol. 1 April 2004: 163–176.

[Abstract](#) [Full Text](#) [Full Text \(PDF\)](#)

Construction cost and invasive potential: comparing *Lythrum salicaria* (Lythraceae) with co-occurring native species along pond banks

Am. J. Bot. 1 December 2001: 2252–2258.

[Abstract](#) [Full Text](#) [Full Text \(PDF\)](#)

Colloquium Paper: Declines of biomes and biotas and the future of evolution

Proc. Natl. Acad. Sci. USA 8 May 2001: 5471–5476.

[Abstract](#) [Full Text](#) [Full Text \(PDF\)](#)

Colloquium Paper: Human–modified ecosystems and future evolution

Proc. Natl. Acad. Sci. USA 8 May 2001: 5458–5465.

[Abstract](#) [Full Text](#) [Full Text \(PDF\)](#)

Impact of Dilution on Microbial Community Structure and Functional Potential: Comparison of Numerical Simulations and Batch Culture Experiments

Appl. Environ. Microbiol. 1 February 2001: 702–712.

[Abstract](#) [Full Text](#) [Full Text \(PDF\)](#)

Environment and health: 7. Species loss and ecosystem disruption -- the implications for human health

CMAJ 9 January 2001: 66–69.

[Full Text](#) [Full Text \(PDF\)](#)

Insect herbivory accelerates nutrient cycling and increases plant production

Proc. Natl. Acad. Sci. USA 19 December 2000: 14412–14417.

[Abstract](#) [Full Text](#) [Full Text \(PDF\)](#)

Assessment of Microbial Diversity in Four Southwestern United States Soils by 16S rRNA Gene Terminal Restriction Fragment Analysis

Appl. Environ. Microbiol. 1 July 2000: 2943–2950.

[Abstract](#) [Full Text](#) [Full Text \(PDF\)](#)

Global Biodiversity Scenarios for the Year 2100

Science 10 March 2000: 1770–1774.

[Abstract](#) [Full Text](#) [Full Text \(PDF\)](#)

Poverty, Defense, and the Environment: How Policy Optics, Policy Incompleteness, fastthinking.com, Equivalency Paradox, Deliberation Trap, Mailbox Dilemma, the Urban Ecosystem, and the End of Problem Solving Recast Difficult Policy Issues

Administration & Society 1 January 2000: 687–725.

[Abstract](#) [Full Text \(PDF\)](#)

Evolutionary consequences of niche construction and their implications for ecology

Proc. Natl. Acad. Sci. USA 31 August 1999: 10242–10247.

[Abstract](#) [Full Text](#) [Full Text \(PDF\)](#)

Biodiversity and ecosystem productivity in a fluctuating environment: The insurance hypothesis

Proc. Natl. Acad. Sci. USA 16 February 1999: 1463–1468.

[Abstract](#) [Full Text](#) [Full Text \(PDF\)](#)

Hypsographic demography: The distribution of human population by altitude

Proc. Natl. Acad. Sci. USA 24 November 1998: 14009–14014.

[Abstract](#) [Full Text](#) [Full Text \(PDF\)](#)

Modulation of Diversity by Grazing and Mowing in Native Tallgrass Prairie

Science 1 May 1998: 745–747.

[Abstract](#) [Full Text](#)

Insect herbivory accelerates nutrient cycling and increases plant production

Proc. Natl. Acad. Sci. USA 19 December 2000: 14412–14417.

[Abstract](#) [Full Text](#) [Full Text \(PDF\)](#)